

Least-Cost Feed Formulation for Juvenile *Macrobrachium rosenbergii* (De Man) by Using the Linear Programming Technique

A.T. LAW, Y. T. POH, and K. J. ANG

Faculty of Fisheries and Marine Science

Universiti Pertanian Malaysia

43400 UPM, Serdang, Selangor Darul Ehsan Malaysia

19278

Keywords: Feed formulation, linear programming, juvenile *Macrobrachium rosenbergii*

ABSTRAK

Pemrograman kos-terkecil linear telah digunakan ke atas juvena *M. rosenbergii* untuk merumuskan satu formula makanan dengan menggunakan bahan-bahan asli (hampas ikan, hampas udang, kelapa kering, hampas kacang soya, tepung gandum dan minyak kelapa sawit). Berikut adalah kriteria yang digunakan: Kandungan asid amino perlu adalah hampir sama dengan kandungan asid amino perlu dalam juvena *M. rosenbergii*, lemak kasar 5-10% dan tenaga kasar 4,400 cal/g dengan kos yang minimum. Empat jenis makanan dihasilkan dengan julat protein di antara 25% hingga 50%. Tindakbalas kadar pertumbuhan juvena *M. rosenbergii* ke atas makanan yang dirumus ini, menunjukkan bahawa makanan yang mempunyai 40% kandungan protein (P40) memberikan kadar pertumbuhan dan nisbah penukaran makanan yang terbaik. Makanan P40 adalah disyorkan sesuai untuk juvena *M. rosenbergii*.

ABSTRACT

Linear least-cost programming was used in juvenile *M. rosenbergii* feed formulation using locally available feed ingredients (fish meal, shrimp meal, copra meal, soybean meal, wheat flour and palm oil). The following constraints were established: the essential amino acid contents were closely similar to those of juvenile *M. rosenbergii*, crude fat 5-10% and gross energy 4,400 cal/g with least cost. Four types of feed were produced with protein ranges from 25% to 50%. Growth responses of juvenile *M. rosenbergii* fed these formulated feeds showed that the 40% protein feed (P40) supported the best specific growth rate and feed conversion ratio. P40 feed is recommended for juvenile *M. rosenbergii*.

INTRODUCTION

Feed costs for *M. rosenbergii* production can account for 30-40% of total operation costs (Liao and Chao 1982). Formulation of a least-cost nutritionally balanced feed would offer an opportunity of reducing overall production cost as well as increasing production. The application of linear programming in least-cost feed formulation for livestock and poultry has gained wide recognition. Its use in fish diet formulation was described by Chow *et al.* (1980). The linear programming technique in feed formulation for *M. rosenbergii* has not yet been reported. There is a paucity of information on the nutritional requirements for prawns including *M. rosenbergii* in the literature (New 1976; Sick and Millikin 1983). Therefore, a study was undertaken to make use of the available information on prawn

nutrition to formulate a least-cost and nutritionally balanced feed from locally available ingredients for *M. rosenbergii* juveniles.

The constraints for the linear programme were set as follows: (1) The amino acid content of the feed was similar to that obtained from the muscle of juvenile *M. rosenbergii*. Many studies have indicated that artificial diets containing essential amino acid profiles that were similar to those of the cultured prawns including *M. rosenbergii* produced better growth (Farmanfarman and Lauterio 1979 and 1980; Kanazawa 1985; Pascual 1989) (2) Crude fat content of the diet was confined between 5 to 10%. Diets which have 10% or more lipid have an adverse effect on the growth and survival of penaeid prawns (Bautista 1986; Sheen and D'Abramo 1991); (3) The gross energy of 4,400

cal/g was based on the reports by Sedgwick (1979); and (4) Based on the wholesale price of the ingredients, the feed production cost was minimal.

MATERIALS AND METHODS

Proximate Analyses

Fish meal, shrimp meal, soybean meal, copra meal and wheat flour were the ingredients used for this study. Their moisture contents, crude fat and crude protein compositions were analyzed according to the methods of A.O.A.C. (1975). Gross energy content was determined by using a bomb calorimeter (Parr Adiabatic Calorimeter). For palm oil, only the crude fat and gross energy contents were determined. *M. rosenbergii* of two different sizes, i.e. 4.5-5.4 cm orbital length (1.35-1.55g) and 6.2-6.8 cm orbital length (1.92-2.20g) and other formulated feeds were also subjected to the same analyses. Six replicates were conducted on each sample.

Amino Acid Analysis

Amino acid contents in the ingredients, juvenile *M. rosenbergii*, and the formulated diets were analyzed following the method of Spitz (1973). A Technicon amino acid analyser (TSM) was used for the analysis with Sigma standard amino acid solution (A.A.S. 18) as a reference. Three replicates were run on each sample.

Linear programming for feed formulation

Four types of feed, i.e. P25 (25% protein), P30 (30% protein), P40 (40% protein), and P50 (50% protein) were formulated. The following constraints were established for the programming: essential amino acid contents were similar to those of the muscle of juvenile *M. rosenbergii*, crude fat 5-10%, gross energy 4,400 cal/g, and least-cost for the ingredients. The wholesale price for fish meal, shrimp meal, soybean meal, copra meal, wheat flour, and palm oil at the time of this study was RM 1.40, RM 1.00, RM 0.88, RM 0.35, RM 0.60, and RM 1.00 (Malaysian Ringgit) per kilogram, respectively. The linear programming was executed using the Functional Mathematical Programming System (FMPS) software package (Sperry Univac). Three per cent of mineral mix, 0.2% vitamin premix, 0.5% basfin (as binder) and 0.25% of calcium propionate (as fungicide) were included for the final feed preparation. The cost of the additives was added

to the least-cost of the individual feed to give the final cost of the prepared feed.

Growth Response of Prawns Fed with the Formulated Feeds

Weight gains were used as the criterion for evaluation of the growth responses of prawns fed with the formulated feeds. The initial weight of prawns used for this study ranged from 0.014g to 0.336g. The prawns were fed at 5% body weight twice a day at 0900 and 2100. Triplicate experiments were conducted on each diet. The water quality parameters of the culture water were as follow: temperature 27-28 °C; dissolved oxygen 6.8-7.8 ppm; pH 6.9-7.3; total ammonium-nitrogen, < 0.02 ppm. Three culture systems were used for the experiments: the aquarium series, basin series and circular tanks. The aquarium series consisted of fifteen glass aquariums measuring 100cm x 60cm x 50cm (length, width and height). Each aquarium was partitioned off 1/3 of its length to form a biological filter section comprising limestone chips and cockle shells with a filtration rate of 6 litres of water per minute. Fifty postlarvae were randomly assigned to each aquarium with adequate shelters as hiding places.

The basin culture system consisted of a series of twelve basins receiving water from the same recirculating system. The flow rate of water into each basin was 500 ml per minute. The dimension of each culture unit was 49cm x 60cm x 30cm (bottom diameter, top diameter and height). The depth of water in each basin was 24 cm. Forty-two and fifteen postlarvae were randomly assigned to each basin in series 1 of 11 experiments respectively. For the circular tanks, the dimension of each unit was 150 cm diameter x 40 cm height (water depth). The culture system consisted of an undergravel filter 10 cm thick and the water flow rate was 1.5 litres per minute.

RESULTS

The proximate analyses of crude protein, fat, ash, moisture, and gross energy of the ingredients and the juvenile *M. rosenbergii* are shown in Table 1. The essential amino acid composition of the ingredients as well as that of the fresh muscle content of the juvenile prawns and the formulated feeds are shown in Tables 2 and 3.

TABLE 1
Proximate analyses of feed ingredients and juvenile *M. rosenbergii*

	Ash (%) ¹	Crude fat (%) ¹	Crude protein (%) ¹	Gross energy (cal/g)	Moisture (%)
Ingredients					
Fish meal	26.90 ± 0.42 ²	9.95 ± 0.07	65.32 ± 0.09	4433.6 ± 43.4	11.91 ± 0.05
Shrimp meal	27.90 ± 0.24	2.26 ± 0.09	42.35 ± 0.81	3843.0 ± 35.2	16.29 ± 0.14
Soybean meal	6.09 ± 0.05	4.97 ± 0.05	45.14 ± 0.28	4808.5 ± 7.2	9.73 ± 0.14
Copra meal	6.28 ± 0.02	13.12 ± 0.15	24.97 ± 0.08	4926.5 ± 12.9	10.19 ± 0.12
Wheat flour	0.45 ± 0.00	1.17 ± 0.03	14.24 ± 1.11	4358.2 ± 14.0	3.11 ± 0.18
Palm oil	- ³	100	-	9480.2 ± 0.0	-
<i>M. rosenbergii</i>					
6.20 - 6.82 cm ⁴ (1.92 - 2.20) ⁵	18.87 ± 0.07	5.84 ± 0.08	63.32 ± 0.33	4609.6 ± 11.3	-
4.50 - 5.35 cm (1.35 - 1.55 g)	20.09 ± 0.09	5.28 ± 0.17	63.59 ± 0.12	4468.0 ± 30.9	-

1: On dry weight basis.

2: Mean ± S.D. (n ± 6).

3: Not determined

4: Post-orbital length.

5: Body weight

TABLE 2
Essential amino acid content of feed ingredients

	Fish meal	Shrimp meal	Soybean meal	Copra meal	Wheat flour
	% Protein (% dry weight)				
Histidine	2.43 ¹ (1.59) ²	2.06(0.93)	3.11(1.40)	4.76(1.19)	5.75(0.82)
Arginine	6.02(3.93)	5.75(2.60)	6.82(3.08)	9.63(2.41)	2.41(0.34)
Valine	4.79(3.25)	4.36(1.97)	4.34(1.96)	3.62(0.90)	3.51(0.50)
Methionine	2.77(1.65)	3.06(1.38)	2.06(0.93)	1.28(0.32)	1.76(0.25)
Isoleucine	3.74(2.44)	2.64(1.19)	3.11(1.40)	2.27(0.57)	2.52(0.36)
Leucine	7.42(4.85)	5.29(2.39)	6.36(2.87)	4.81(1.20)	5.53(0.79)
Tyrosine	2.64(1.72)	7.29(3.29)	3.80(1.72)	1.29(0.32)	0.93(0.13)
Phenylalanine	3.65(2.38)	3.89(1.76)	5.35(2.41)	3.57(0.89)	3.93(0.56)
Lysine	7.82(5.11)	8.07(3.65)	6.67(3.01)	7.41(1.85)	10.85(1.55)
Threonine	4.58(2.99)	4.01(1.81)	3.41(1.54)	2.48(0.62)	2.35(0.33)

1: Calculated from % of essential amino acid on dry weight basis x (1/% crude protein on dry weight basis).
2: Mean value from three analyses, on dry weight basis.

TABLE 3
Essential amino acid content of the formulated feeds and juvenile
M. rosenbergii (1.55g - 2.20g)

	<i>M. rosenbergii</i>	P25	P30	P40	P50
	% Protein (% dry weight)				
Histidine	1.61 ¹ (1.07) ²	1.80(0.45)	2.31(0.69)	1.40(0.56)	1.23(0.61)
Arginine	3.84(2.55)	4.24(1.06)	4.63(1.39)	3.90(1.56)	3.72(1.86)
Valine	3.11(2.06)	3.20(0.80)	3.10(0.93)	3.09(1.24)	3.08(1.54)
Methionine	2.91(1.56)	2.66(0.67)	3.42(1.04)	2.06(0.81)	1.79(0.89)
Isoleucine	2.92(1.94)	2.92(0.73)	2.57(0.77)	2.69(1.08)	2.78(1.39)
Leucine	5.05(3.34)	6.52(1.63)	5.70(1.71)	6.05(2.42)	5.61(2.80)
Tyrosine	3.87(2.57)	1.48(0.37)	1.23(0.37)	1.63(0.65)	1.76(0.88)
Phenylalanine	3.04(2.02)	4.24(1.06)	3.73(1.12)	3.64(1.46)	3.13(1.56)
Lysine	6.41(4.26)	4.24(1.06)	4.10(1.23)	3.83(1.53)	4.50(2.25)
Threonine	2.50(1.66)	2.92(0.73)	2.03(0.61)	3.24(1.30)	2.72(1.36)

1: Calculated from % of essential amino acid on dry weight basis x (1/% crude protein on dry weight basis)
2: Mean value from three analyses

respectively. The ingredient composition, proximate analyses and cost of the formulated feeds are presented in Table 4. The growth responses of *M. rosenbergii* postlarvae and juveniles fed with the formulated feeds in the different culture systems are given in Table 5. The apparent digestibility of the feeds is given in Table 6.

DISCUSSION

Many studies have shown that prawns including *M. rosenbergii* require dietary sources of essential amino acids for growth (Watanabe 1975; NRC

1983; Kanazawa 1985). Depressed appetite and reduced growth rate of the prawns are usually reported when the prawns are fed with diets deficient in any of the essential amino acids. Farmanfarmaian and Lauterio (1979, 1980) reported that the Purina Marine Ration (M20) was insufficient in some of the essential amino acids such as isoleucine, lysine and arginine for juvenile *M. rosenbergii*. However, when it was adjusted to the amino acid pattern of the prawn, it became the effective source of protein. This clearly supported the phenomenon that the

TABLE 4
Ingredient composition, proximate analyses and cost of the formulated feeds

Ingredients, kg/kg of feed.	Formulated feeds			
	P25	P30	P40	P50
Fish meal	0.096	0.116	0.154	0.456
Shrimp meal	0.200	0.224	0.439	0.309
Soybean meal	0.050	0.200	0.100	0.100
Copra cake	0.010	0.114	0.150	0.013
Wheat flour	0.503	0.346	0.133	0.100
Palm oil	0.076	0.050	0.025	0.023
Fine sand	0.065	0.050	0.000	0.000
Proximate analyses (on dry weight basis)				
Crude protein (%)	25.37 ± 0.12 ¹	30.34 ± 0.47	39.50 ± 0.07	49.48 ± 0.41
Crude fat (%)	8.09 ± 0.09	8.35 ± 0.09	6.91 ± 0.14	8.07 ± 0.03
Ash (%)	18.15 ± 0.11	17.52 ± 0.03	19.64 ± 0.00	22.03 ± 0.06
Gross energy (cal/g)	4283.6 ± 0.5	4229.3 ± 46.2	4206.7 ± 39.2	4344.1 ± 5.5
Cost with additives (RM/kg) ²	0.94	0.95	1.08	1.30

1: Mean ± S.D. (n = 6). 2: US \$1 ± RM 2.5 (Malaysian Ringgit)

closer the amino acid composition of the feed to that of the prawn, the more effective is the protein source. Therefore, based on this phenomenon, four types of feeds with protein levels ranging between 25 % to 50 % where the amino acid content was similar to that of the whole fresh muscle of juvenile *M. rosenbergii* were formulated.

The nutrient contents of the formulated feeds were remarkably close to the constraints set by the linear programme (Table 4). The feeds were isocaloric with gross energy content of about 4,400 cal/g. The amino acid contents in the feeds, in terms of per cent protein, were very close to the values obtained from the fresh muscle amino acid content of juvenile *M. rosenbergii*, except for tyrosine which was about 50% lower. The reason for lower tyrosine levels in the feed is not known. Most probably, it was lost during the feed preparation. The water stability of the feeds was greater than 3 hours at 28°C.

The performance of the feeds was evaluated by using the growth responses and feed conversion ratio of postlarvae and juvenile *M. rosenbergii* fed with these feeds in the aquarium and basin culture systems for a period of 33-61 days. The results indicated that under both culture systems, all formulated diets produced

specific growth rates that were significantly different ($P < 0.01$) where P40 feed supported the highest growth rate followed by P50, P30, and P25 feeds. In general, P40 feed also showed better feed conversion ratio (1.68-2.03) among all the feeds, except for P30 feed which was comparable or sometimes better (Table 5). The decline in growth rate and feed conversion ratio of P50 feed suggested that the optimal protein level in the diet for *M. rosenbergii* postlarvae and juveniles was around 40% with a gross energy of 4,400 cal/g, a protein:gross energy ratio of 1:110 and an amino acid profile similar to that of the prawn. The cost of ingredients required for the production of 1 kilogram of P40 feed was RM 1.08 (US\$1.00 = RM 2.50).

It is difficult to compare the results of this study with those of other studies on the growth of juvenile *M. rosenbergii* on the pelleted feed because of the diverse culture systems, stocking densities, and different protein levels in the feed. However, it is interesting to note that the feeds used in this study seemed to give a higher specific growth rate and conversion ratio as compared to the studies conducted elsewhere with feeds of the same protein level, and under similar culture conditions. The better performance of the feeds used in the study may be due to the

TABLE 5
Growth responses of postlarvae and juvenile *Macrobrachium rosenbergii* fed on the formulated diets

Culture system	Type of feed	Stocking density (no/unit)	Initial weight (g)	Final weight (g)	Culture period (days)	Specific growth rate (% per day)	FCR	Survival rate (%)
Aquarium series	P50 (3) ¹	50 (111) ²	0.014	0.541 ± 0.0393	61	5.99 ± 0.12	2.13 ± 0.03	80.67 ± 5.03
	P40 (3)	50 (111)	0.014	0.652 ± 0.017	61	6.30 ± 0.04	2.03 ± 0.09	82.00 ± 2.00
	P30 (3)	50 (111)	0.014	0.487 ± 0.18	61	5.82 ± 0.06	1.98 ± 0.05	82.67 ± 4.16
	P25 (3)	50 (111)	0.014	0.338 ± 0.22	61	5.22 ± 0.11	2.39 ± 0.24	83.33 ± 3.06
	RBS (3)	50 (111)	0.014	0.300 ± 0.028	61	5.02 ± 0.11	2.55 ± 0.06	80.67 ± 9.02
Basin series-I	P50 (3)	42 (221)	0.104 ± 0.006	0.347 ± 0.036	35	3.43 ± 0.22	1.89 ± 0.09	90.47 ± 8.60
	P40 (3)	42 (221)	0.103 ± 0.004	0.389 ± 0.024	35	3.66 ± 0.17	1.86 ± 0.07	82.80 ± 4.10
	P30 (3)	42 (221)	0.108 ± 0.011	0.327 ± 0.034	35	3.16 ± 0.18	1.81 ± 0.19	93.80 ± 4.42
	P25 (3)	42 (221)	0.100 ± 0.006	0.263 ± 0.034	35	2.72 ± 0.20	2.48 ± 0.16	83.30 ± 9.80
Basin series-II	P50 (3)	15 (79)	0.347 ± 0.008	1.842 ± 0.054	58	2.88 ± 0.07	1.89 ± 0.45	84.47 ± 3.87
	P40 (3)	15 (79)	0.336 ± 0.015	2.072 ± 0.169	58	3.13 ± 0.11	1.68 ± 0.02	82.23 ± 3.87
	P30 (3)	15 (79)	0.342 ± 0.013	1.543 ± 0.061	58	2.60 ± 0.02	1.67 ± 0.02	86.70 ± 0.00
	P25 (3)	15 (79)	0.345 ± 0.003	1.117 ± 0.030	58	2.11 ± 0.05	2.43 ± 0.07	80.00 ± 0.00
Circular Tank	P30 (1)	50 (28)	1.590 ± 0.050	4.822 ± 0.250	50	2.70	1.78	90.00
	P30 (1)	100 (56)	1.35		15.05		159	1.52

1: Parenthesis indicates the number of replicates

2: Number of prawns per unit of culture system (Number of prawns per m²); 3: Mean ± S.D.

TABLE 6
Apparent nutrient digestibility of P30 and P40 feeds by juvenile (0.495 g-1.271g) and adult (7.954 g-25.588 g) *M. rosenbergii*

	Apparent digestibility (%) ¹					
	Protein	Fat	Carbohydrate	Ash	Gross energy	Dry matter
P30 Feed						
Juvenile	66.02 ± 4.25 ²	83.48 ± 6.11	82.16 ± 1.83	27.27 ± 9.37	78.77 ± 6.03	62.83 ± 3.84
Adult	72.59 ± 2.37	84.12 ± 2.86	72.15 ± 4.07	41.34 ± 12.61	70.89 ± 2.66	64.94 ± 2.15
P40 Feed						
Juvenile	66.50 ± 3.30	79.61 ± 0.69	63.81 ± 6.86	33.27 ± 4.49	65.77 ± 3.67	57.17 ± 3.23
Adult	75.93 ± 0.36	85.78 ± 0.51	70.35 ± 4.50	43.10 ± 3.47	74.44 ± 3.19	65.77 ± 1.13

1: On dry weight basis. 2: Mean ± S.D. (n = 6)

amino acid pattern in the feed which was similar to that of the muscle of the prawn. A specific growth rate of 5.22 % per day in the aquarium experiment using P25 feed was comparable to that of a 25% protein feed as reported by Willis *et al.* (1976). They obtained a specific growth rate of 5.71 % per day and a feed conversion ratio of 5.50, while we obtained a similar growth rate but a much better feed conversion ratio of 2.39.

Millikin *et al.* (1980) reported a specific growth rate of 2.59% per day for *M. rosenbergii* fed on a 40% protein feed and grown in basin. Using the similar culture system, we obtained a specific growth rate of 3.80 % per day with the P40 feed. In the basin experiment, our P25 feed produced a much better feed conversion ratio of 2.43 compared to that of 3.16-8.16 for 25% protein feed reported by Sandifer and Smith (1977). Even among the treatments in the aquarium experiments of this study, P25 feed produced a comparable higher specific growth rate and a better feed conversion ratio (5.22 % per day & 2.39) than the repelleted broiler starter (RBS) which had the same levels of protein, fat and gross energy (5.02% & 2.55).

In the circular tank experiments, the juvenile *M. rosenbergii* weighing 1.35 g achieved an average weight of 15.05 g after 159 days when fed with the P30 feed. The specific growth rate was 1.52 % per day and the feed conversion ratio was 2.55. These values were better than those obtained from juvenile *M. rosenbergii* that were fed with the Purina Marine 25 which was supplemented with fresh beef liver; the specific rate and conversion ratio were 0.81 % per day

and 5.18 respectively (Farmanfarmaian and Lauterio 1979). These results seem to support the phenomenon that feeds containing protein having a similar amino acid pattern to the culture species will be a more effective source of protein for growth.

The findings of this study revealed that the 40 % protein formulated feed (P40) supported the optimal growth of *Macrobrachium rosenbergii*. In fact, many studies have also indicated that the optimal protein requirement for *M. rosenbergii* is about 40 % (Pandian 1989). Millikin *et al.* (1980) demonstrated that juvenile *M. rosenbergii* fed with a 40% protein diet had a better weight gain than those fed with 49%, 32% and 23% protein diets over a period of 14 weeks. The results of the protein requirement studies of juvenile *M. rosenbergii* by Chao (1979) suggested that a protein level in excess of 35% was required for optimum growth of the prawn. Willis *et al.* (1976) showed that a 40% protein diet (trout chow) produced better growth than the 25% protein diet of Purina Marine Ration 25. Natural productivity in the pond plays an important role in providing supplemental diets for the growth of *M. rosenbergii* fed on pelleted feed (Fair and Fortner 1981). Therefore, in practice, farmers found that pelleted feeds containing 15-25% protein were satisfactory for the culture of *M. rosenbergii* in earthen ponds that had been fertilized with manure (New 1976; New 1990). Feed cost accounts for a high percentage of the production cost for *M. rosenbergii*. To reduce the cost of production, P30 feed has been successfully used along with chicken manure fertilization of the pond for *M. rosenbergii* production (Maclean *et al.* 1989).

ACKNOWLEDGEMENTS

This research was supported by grants from Universiti Pertanian Malaysia and EEC No.TSD A 287 Mal (H).

REFERENCES

- ASSOCIATION OF OFFICIAL ANALYTICAL CHEMISTS. 1975. *Official Methods of Analysis*. Washington, D.C.: A.O.A.C. 1094 pp.
- BAUTISTA, M.N. 1986. The response of *Penaeus monodon* juveniles to varying protein/energy ratios in test diets. *Aquaculture* **53**: 229-242.
- CHAO, N.H. 1979. Freshwater prawn farming in Taiwan - the patterns, problems and prospects. *Proc. World Maricult. Soc.* **10**: 51-67.
- CHOW, K.W., G. L. RUMSEY and P. W. WALDROUP. 1980. Linear programming in fish diet formulation. In: *Fish Feed Technology*. ADCP/REP/80/11, p. 241-286.
- FAIR, P.H. and A. R. FORTNER. 1981. The role of formula feeds and natural productivity in culture of the prawn, *Macrobrachium rosenbergii*. *Aquaculture* **24**: 233-243.
- FARMANFARMAIAN, A. and T. LAUTERIO. 1979. Amino acid supplementation of feed pellets of the giant shrimp *M. rosenbergii*. *Proc. World Maricult. Soc.* **10**: 674-688.
- FARMANFARMAIAN, A. and T. LAUTERIO. 1980. Amino acid composition of the tail muscle of *M. rosenbergii* - comparison to amino acid pattern of supplemented commercial feed pellets. *Proc. World Maricult. Soc.* **11**: 454-462.
- KANAZAWA, A. 1985. Nutrition of Penaeid prawns and shrimps. In Y. Taki, J.H. Primavera, J.A. Llobrera (eds.), *Proc. 1st. Intl. Conf. on culture of Penaeid Prawns/Shrimps*. Iloilo City, Philippines. Aquaculture Dept. SEAFDEC, p. 123-130.
- LIAO, I-C. and N. H. CHAO. 1982. Progress made in *M. rosenbergii* farming and its extension in Taiwan. In M.B. New (ed.), *Giant Prawn Farming*. Amsterdam: Elsevier. p. 357-381.
- MACLEAN, M. H., K. J. ANG, J. H. BROWN and K. JAUNCEY 1989. The effect of organic fertilizer and formulated feed in pond culture of the freshwater prawn, *Macrobrachium rosenbergii* (de Man): prawn production. *Aquacult. Fish. Manage.* **20**: 399-406.
- MILLIKIN, M.R., A. R. FORTNER and P. H. FAIR. 1980. Influence of dietary protein concentration on growth, feed conversion and general metabolism of juvenile prawn (*Macrobrachium rosenbergii*). *Proc. World Maricult. Soc.* **11**: 382-391.
- NATIONAL RESEARCH COUNCIL. 1983. Nutrient Requirements of Warmwater Fishes and Shellfishes. Revised Edition. Washington, D.C.: National Academic Press. 102 pp.
- NEW, M.B. 1976. A review of dietary studies with shrimp and prawns. *Aquaculture* **9**: 101-144.
- NEW, M.B. 1990. Freshwater prawn culture: a review. *Aquaculture* **88**: 99-143.
- PANDIAN, T.J. 1989. Protein requirements of fish and prawns culture in Asia. In: S.S. De Silva, (ed.), *Fish Nutrition Research in Asia. Proceedings of the Third Asian Fish Nutrition Network Meeting*. Asian Fish. Soc. Spec. Publ. **4**. 166 pp. Manila, Philippines: Asian Fisheries Society.
- PASCUAL, F.P. 1989. Status of shrimp nutrition and feed development in Southeast Asia. In S.S. De Silva (ed.), *Fish Nutrition Research in Asia. Proceedings of the Third Asian Fish Nutrition Network Meeting*. Asian Fish. Soc. Spec. Publ. **4**. 166 pp. Manila, Philippines: Asian Fisheries Society.
- SANDIFER, P.A. and T.I.J. SMITH. 1977. Intensive rearing of postlarval Malaysian prawns, *M. rosenbergii* in a closed cycled nursery system. *Proc. World Maricult. Soc.* **8**: 235-255.
- SEDGWICK, R. 1979. Influence of dietary protein and energy on growth, food consumption and food conversion efficiency in *P. merguensis*. *Aquaculture* **16**: 7-10.
- SHEEN, S.-S. and L.R. D'ABRAMO. 1991. Response of juvenile freshwater prawn, *Macrobrachium rosenbergii*, to different levels of a cod liver oil/corn oil mixture in a semi-purified diet. *Aquaculture* **93**: 121-134.
- SICK, L.V. and M. R. MILLIKIN. 1983. Dietary and nutrient requirements for culture of the Asian prawn, *Macrobrachium rosenbergii*. In: J.P. McVey (ed.), *CRC Handbook of Mariculture. Vol. I. Crustacean Aquaculture*. Boca Raton, FL: CRC Press, Inc. p. 381-389.
- SPITZ, H.D. 1973. A new approach for sample preparation of protein hydrolysates for amino

acid analysis. *Anal. Biochem.* **56**: 66-73.

WATANABE, W.O. 1975. Identification of essential amino acids of the fresh-water prawn, *Macrobrachium rosenbergii*. Masters thesis. University of Hawaii. Honolulu. Hawaii, USA.

WILLIS, S.A., R. W. HAGOOD and G. T. ELIASON. 1976. Effects of four stocking densities and three diets on growth and survival of postlarval *M. rosenbergii* and *M. acanthurus*. *Proc. World Maricult. Soc.* **7**: 655-662.

(Received 3 May 1993)

ABSTRACT

SUMMARY