

Selection of Broodstock of Tiger Prawn, *Penaeus monodon* Fabricius, on the Basis of Morphometric Traits

S.K. DAUD and K. J. ANG

Faculty of Fisheries and Marine Sciences

Universiti Pertanian Malaysia

43400 UPM, Serdang, Selangor, Malaysia

Keywords: broodstock selection, tiger prawn, *Penaeus monodon*, morphometric traits

ABSTRAK

Penaeus monodon matang dipungut dari perairan Kedah, Perak, Terengganu, Johor dan Sabah. Ciri morfometrik terpilih dan berat badan bagi setiap ekor udang disukat. Udang betina didapati lebih besar ($P < 0.05$) berbanding dengan udang jantan bagi semua lokasi. Analisis regresi langkah demi langkah menunjukkan bahawa panjang karapas sahaja mencukupi untuk menerangkan varians dalam berat badan bagi udang jantan dan betina. Panjang karapas boleh memberikan anggaran yang lebih tepat bagi berat badan dan oleh itu ia boleh dijadikan sebagai satu kriteria dalam pemilihan induk. Hubungan logaritma antara berat badan dan panjang penuh, berat badan dan panjang karapas, dan antara panjang karapas dan panjang penuh bagi kedua-dua seks juga dikira.

ABSTRACT

Mature *Penaeus monodon* were collected from Kedah, Perak, Terengganu, Johore and Sabah waters in Malaysia. Selected morphometric characters and body weight were measured individually. The results showed that the females were significantly larger ($P < 0.05$) than the males in all locations. Stepwise regression analysis indicated that carapace length alone is sufficient to explain the variance in body weight for male and female prawns and can thus be used as a criterion for the selection of broodstock. The logarithmic relationships between body weight and total length, body weight and carapace length, and between carapace length and total length for both sexes were also calculated.

INTRODUCTION

Penaeus monodon Fabricius, commonly known as tiger prawn, is the main cultured prawn species in Malaysia. Despite the establishment of hatchery and rearing techniques for this species, prawn broodstocks used for seed production and selective breeding are still being collected from natural populations.

Morphometric traits can easily be used as a criterion for the selection of prawn broodstock (Lester 1983; Huang *et al.* 1990) without killing or damaging them. Total length and carapace length are commonly used to estimate the size composition of prawn species (Ramamurthy and Manickaraja 1978). In addition, some morphometric characters can give a more accu-

rate estimate of body weight and can be used for selecting broodstock on the basis of weight. Relationships between total length and carapace length, and total length and weight have been described for *P. monodon* and other prawn species (Hall 1962; Kirkegaard *et al.* 1970; Ramamurthy and Manickaraja 1978; Horton and Lester 1982; Lester 1983; Goswami *et al.* 1986; Huner *et al.* 1988).

The objectives of this study were to estimate the variability of selected morphometric traits and to determine the relationships among the characters in tiger prawns from different localities.

MATERIALS AND METHODS

Samples of wild and mature *Penaeus monodon* were purchased alive from fishermen in five

different areas, Kedah, Perak, Terengganu, Johore and Sabah waters.

Individual prawns were weighed in the laboratory to the nearest 0.001 g. Total length and carapace length were measured using vernier calipers to the nearest 0.01 cm. All parameters were based on Chua (1978), Grey *et al.* (1983), and Ahmad and Mahyam (1986). The proportion of carapace length to total length or carapace length index (%) was calculated by (carapace length/total length) x 100. Differences in carapace length indices among localities and between sexes were tested by one-way analysis of variance. Tukey test analysis was used to measure differences in carapace length index mean values among locations. The characters were log transformed to reduce the correlation of means and variances (Sokal and Rohlf 1981).

The morphometric characters were analysed by stepwise variable selection regression method using Statgraphics statistical graphics system (STSC, Inc. and Statistical Graphics Corporation, USA). Body weight was used as a dependent variable while total length and carapace length were independent variables. The relationships between total length and weight, and between carapace length and weight are described by the allometric equation, i.e. $Y = aX^b$, where X and Y are independent and dependent variables, respectively, and a and b are two constants. Non-linear regression analysis (Sokal and Rohlf 1981) was performed to determine these relationships for both sexes from different locations. Linear regression analysis was also used to describe the relationship between carapace length and total length. Data were pooled for males and females to provide a common regression equation describing the relationships between the two variables.

RESULTS AND DISCUSSION

The range, mean and coefficient of variation values for adult male and adult female shrimps collected from the five locations are presented in Table 1. Females were significantly larger ($P < 0.05$) than the males, with average body weights of 113.54 g and 74.88 g, respectively. This characteristic was also noted in *Penaeus longistylus* (Penn 1980), *P. latissulcatus* and *P. esculentus* Haswell (Penn and Hall 1974). Male shrimps, however, had less variable and greater carapace length indices than the females, indicating that carapace length increases more rapidly than the

total length in males but less rapidly in females. The coefficient of variation of the carapace length indices ranged between 5 and 18% in males and 5 to 21% in females. Among the various locations, males from Sabah had the highest average carapace length index (36.46%) and males from Kedah the lowest (22.65%). Similarly to males, the carapace length index (%) of females from Sabah was the largest (36.43%) and females from Kedah were also the smallest (23.17%). The Tukey test shows that the carapace length indices of females were significantly different ($P < 0.05$) among locations, except between those from Johore and Perak. The prawns from Johore and Perak could be derived from the same population. The difference in size of carapace within sexes among different locations may be due to different stocks and different environments (Goswami *et al.* 1986; Huang *et al.* 1990).

The results of stepwise variable selection regression analysis for male and female prawns from five locations are shown in Table 2. In males, the carapace length was the first variable in stepwise regression analysis in Perak and Kedah populations, accounting for 79.4% and 72.4%, respectively, of the variation in body weight. Total length became the first entered variable in the regression analysis in Sabah, Terengganu and Johore male shrimp populations, explaining 77.8%, 61.8% and 61.0% of the variation in body weight, respectively. Conversely, among females, carapace length was the first variable to enter the regression step in Perak and Johore populations, explaining 95.6% and 79.7% of the variations in body weight, respectively. The total length was the first entered variable in stepwise regressions of Sabah, Kedah and Terengganu female populations that explained 82.88%, 80.6% and 74.7%, respectively, of the variance in body weight.

This study indicates that carapace length alone is sufficient to explain the variance in body weight for male (65.4%) and female (73.8%) prawns although a combination of two or three variables at a time gives a more accurate estimation of body weight.

From the analysis of variance, there were significant differences ($P < 0.05$) in length between sexes for all locations. Therefore, the regression analyses for males and females were done separately.

Table 3 describes the logarithmic relationships between body weight (W) and total length

TABLE 1

Range, mean and coefficient of variation (CV) of total length (TL), carapace length (CL), body weight (WT) and carapace length index (CL. I.) for male (M) and female (F) *Penaeus monodon* collected from different regions in Malaysia

Variable	Sex	Kedah			Perak			Terengganu			Johore			Sabah		
		Range	Mean	CV	Range	Mean	CV	Range	Mean	CV	Range	Mean	CV	Range	Mean	CV
		M (N=46) F (N=33)			M (N=95) F (N=6)			M (N=89) F (N=65)			M (N=85) F (N=10)			M (N=68) F (N=33)		
TL (cm)	Male	16.2-25.8	20.73	0.12	18.2-28.9	23.34	0.10	15.3-36.0	24.42	0.12	18.8-26.2	22.84	0.07	18.1-26.3	22.48	0.07
	Female	15.6-27.7	23.20	0.13	23.5-27.5	20.73	0.06	17.3-36.8	28.37	0.14	22.3-26.1	23.89	0.05	20.5-27.0	23.63	0.07
CL (cm)	Male	3.5-5.6	4.68	0.10	6.7-10.1	8.09	0.08	4.0-9.5	7.02	0.20	6.1-8.8	7.41	0.07	6.6-10.0	8.18	0.09
	Female	4.0-6.8	5.36	0.12	7.3-8.7	8.27	0.07	3.7-10.9	7.41	0.18	6.7-8.5	7.67	0.07	6.3-10.3	8.59	0.09
WT (g)	Male	29.5-104.1	57.01	0.28	47.9-130.4	72.70	0.23	37.5-261.2	79.80	0.40	41.7-90.7	62.12	0.17	52.6-137.8	86.48	0.23
	Female	41.5-132.4	81.92	0.26	55.2-101.4	85.05	0.40	49.1-265.1	133.74	0.40	52.9-83.8	65.94	0.15	62.2-143.5	99.57	0.21
CL. I. (5)	Male	19.2-27.9	22.57	0.09	27.8-40.1	34.65	0.06	20.1-39.2	28.76	0.18	29.0-37.0	32.43	0.05	32.1-41.1	36.40	0.06
	Female	19.9-27.9	23.10	0.08	27.0-34.0	31.64	0.08	12.2-35.9	26.13	0.21	28.6-35.0	32.11	0.05	30.7-40.7	36.38	0.05

TABLE 2

Range, mean and coefficient of variation (CV) of total length (TL), carapace length (CL), body weight (WT) and carapace length index (CL. I.) for male (M) and female (F) *Penaeus monodon* collected from different regions in Malaysia

Variable	Sex	Kedah		Perak		Terengganu		Johore		Sabah	
		b	R	b	R ²	b	R ²	b	R ²	b	R ²
log CL	Male	2.422	0.724	2.511	0.794	1.575	0.501	1.748	0.584	1.941	0.584
	Female	1.833	0.771	3.356	0.956	2.028	0.660	1.756	0.797	1.775	0.673
	Combined	2.260	0.798	2.568	0.794	1.931	0.628	1.748	0.604	1.937	0.632
log TL	Male	1.948	0.691	1.643	0.573	1.986	0.618	1.849	0.610	2.987	0.778
	Female	1.781	0.806	0.409	0.011	2.799	0.747	1.139	0.623	2.738	0.828
	Combined	2.090	0.768	1.604	0.547	2.434	0.689	1.846	0.613	2.891	0.809

TABLE 3
Relations of log body weight (W) to log total length (L) in male and female of *Penaeus monodon* from different locations

Location	Sex	log W = log a + b log L	Correlation coefficient (r)
Kedah	Male	log W = -0.809 + 1.948 log L	0.83
	Female	log W = -0.518 + 1.781 log L	0.90
Perak	Male	log W = -0.418 + 2.511 log L	0.89
	Female	log W = -1.149 + 3.356 log L	0.98
Terengganu	Male	log W = -0.869 + 1.986 log L	0.79
	Female	log W = -1.959 + 2.799 log L	0.86
Johore	Male	log W = -0.719 + 1.849 log L	0.78
	Female	log W = -1.128 + 2.139 log L	0.79
Sabah	Male	log W = -2.101 + 2.987 log L	0.90
	Female	log W = -1.763 + 2.738 log L	0.91
Combined	Male	log W = -0.671 + 1.899 log L	0.62
	Female	log W = -0.513 + 1.741 log L	0.64

(L) in male and female prawns. Non-linear regression equations of $\log W = \log a + b \log L$ are shown for each location (Table 3). The relationships between body weight and total length for males and females in all locations were estimated as $\log W = -0.513 + 1.741 \log L$ and $\log W = -0.671 + 1.899 \log L$, respectively. The values of coefficient "b" of the weight-total length relationships show considerable variation, ranging from 1.899 to 2.987 in males and from 1.741 to 3.356 in females. The "b" value was significantly smaller ($P < 0.05$) than 3.0 for males. In contrast, "b" value in *P. duorarum* was significantly greater than 3.0 for both sexes as reported by Kutkuhn (1962). However, Fontaine and Neal (1971) found that the "b" value for *P. duorarum* males was not different from 3.0 ($P > 0.05$).

The logarithmic relationships between body weight (W) and carapace length (L_{car}) for both males and females from each location are also described by the non-linear regression equation of $\log W = \log a + b \log L_{car}$ (Table 4). For all locations, the relationships between body weight and carapace length for males and females were estimated as $\log W = -0.019 + 2.10 \log L_{car}$ and $\log W = -0.083 + 2.213 \log L_{car}$, respectively.

Table 5 shows the logarithmic relationships between carapace length (L_{car}) and total length

(L) for male and female prawns from each location as $\log L_{car} = \log a + b \log L$. For all locations, the linear relationships between carapace length and total length for males and females were described as $L_{car} = -0.561 + 1.042 \log L$, respectively.

The results show that carapace length increased linearly with the total length although carapace length does not grow allometrically with body weight. However, body weight is correlated more with carapace length for both male and female prawns, $r = 0.81$ and $r = 0.86$, respectively (Table 4) compared with total length with $r = 0.64$ and $r = 0.62$ for males and females, respectively (Table 3). Carapace length is a considerably more stable size reference dimension (Rhodes and Holdich 1984) and this may contribute to its good correlation with body weight. Total length measurement is rather varied according to the length of the rostrum, which is easily damaged, thus affecting the accuracy of this measurement.

This study indicates that carapace length can be used as a single variable to explain a large proportion of the variance in body weight. This is in accordance with the measurement used by Hall (1962). Thus this variable can be regarded as an important criterion for choosing

TABLE 4
Relationship of log body weight (W) to log carapace length (L_{car}) in male and female of
Penaeus monodon from different locations

Location	Sex	$\log W = \log a + b \log L$	Correlation coefficient (r)
Kedah	Male	$\log W = 0.133 + 2.422 \log L_{car}$	0.85
	Female	$\log W = 0.577 + 1.833 \log L_{car}$	0.88
Perak	Male	$\log W = 0.418 + 2.511 \log L_{car}$	0.89
	Female	$\log W = 1.149 + 3.356 \log L_{car}$	0.98
Terengganu	Male	$\log W = 0.471 + 1.575 \log L_{car}$	0.71
	Female	$\log W = 0.101 + 2.028 \log L_{car}$	0.81
Johore	Male	$\log W = 0.273 + 1.748 \log L_{car}$	0.76
	Female	$\log W = 0.265 + 1.756 \log L_{car}$	0.89
Sabah	Male	$\log W = 0.165 + 1.941 \log L_{car}$	0.76
	Female	$\log W = 0.340 + 1.775 \log L_{car}$	0.82
Combined	Male	$\log W = 0.083 + 2.213 \log L_{car}$	0.86
	Female	$\log W = 0.019 + 2.100 \log L_{car}$	0.81

TABLE 5
Relationship of log carapace length (L_{car}) to log total length (L) in male and female of *Penaeus monodon*
from different locations

Location	Sex	$\log L_{car} = \log a + b \log L$	Correlation coefficient (r)
Kedah	Male	$\log L_{car} = -0.110 + 0.593 \log L$	0.72
	Female	$\log L_{car} = -0.329 + 0.775 \log L$	0.82
Perak	Male	$\log L_{car} = 0.111 + 0.583 \log L$	0.76
	Female	$\log L_{car} = 0.741 + 0.124 \log L$	0.11
Terengganu	Male	$\log L_{car} = -0.476 + 0.953 \log L$	0.51
	Female	$\log L_{car} = 0.101 + 2.029 \log L$	0.81
Johore	Male	$\log L_{car} = -0.164 + 0.761 \log L$	0.74
	Female	$\log L_{car} = -0.388 + 0.924 \log L$	0.67
Sabah	Male	$\log L_{car} = 0.486 + 1.035 \log L$	0.78
	Female	$\log L_{car} = -0.700 + 1.190 \log L$	0.86
Combined	Male	$\log L_{car} = 0.084 + 0.548 \log L$	0.37
	Female	$\log L_{car} = -0.561 + 1.042 \log L$	0.53

broodstock in selective breeding programmes of *P. monodon*.

ACKNOWLEDGEMENTS

The authors wish to thank Universiti Pertanian Malaysia for the use of facilities and the Malaysian Government for providing research grant IRPA 1-07-05-073 (50367). Thanks are also due to Fauziah Ahmad and Noriza Harun for their technical assistance.

REFERENCES

- AHMAD, A.T. and M.I. MAHYAM. 1986. Panduan Mengenal-pasti Jenis-jenis Udang Laut Dagangan di Malaysia. Kuala Lumpur: Department of Fisheries, Ministry of Agriculture.
- CHUA, C.W. 1978. *Commercial Prawns of Peninsular Malaysia*. Kuala Lumpur, Department of Fisheries, Ministry of Agriculture.
- FONTAINE, C.T. and R.A. NEAL. 1971. Length-weight relations for three commercially important Penaeid shrimp of the Gulf of Mexico. *Trans. Amer. Fish. Soc.* **3**: 584-586.
- GOSWAMI, U., S.G. DALAL and S.C. GOSWAMI. 1986. Preliminary studies on prawn, *Penaeus merguensis*, for selection of broodstock in genetic improvement programs. *Aquaculture* **53**: 41-48.
- GREY, D.L., W. DALL and A. BAKER. 1983. *A Guide to the Australian Penaeid Prawns*. Darwin: Northern Territory Department of Primary Production.
- HALL, D.N.F. 1962. *Observations on the Taxonomy and Biology of Some Indo-West-Pacific Penaeidae (Crustacea, Decapoda)*. Col. Off. Fish. Publ. No. 17. London, HMSO.
- HORTON, S.E. and L.J. LESTER. 1982. Utilizing morphometric and population genetic data in domestication of Penaeid shrimp. *Proc. World Maricult. Soc.* **13**.
- HUANG, C.M., M.Y. TSAI and I.C. LIAO. 1990. Morphometric studies on choice of broodstock for genetic improvement in *Penaeus penicillatus*. In: *Proceedings Second Asian Fisheries Forum*, ed. R. Hirano and I. Hanyu, p. 523-526. Manila: The Asian Fisheries Society.
- HUNER, J.V., O.V. LINDQVIST and H. KONONEN. 1988. Comparison of morphology and edible tissues of two important commercial crayfishes, the noble crayfish, *Astacus astacus* Linne, and the red swamp crayfish, *Procambarus clarkii* (Girard) (Decapoda, Astacidae and Cambaridae). *Aquaculture* **68**: 45-57.
- KIRKEGAARD, I., D.J. TUMA and R.H. WALKER. 1970. Synopsis of biological data on the banana prawn *Penaeus merguensis* de Man, 1888. *Fisheries Synopsis, No. 8*. Melbourne, CSIRO Fisheries and Oceanography.
- KUTRUHN, J.H. 1962. Dynamics of a Penaeid shrimp population and management implications. *U.S. Fish Wildl. Serv. Fish. Bull.* **65**(2): 313-338.
- LESTER, L.J. 1983. Developing a selective breeding program for Penaeid shrimp mariculture. *Aquaculture* **33**: 41-50.
- PENN, J.W. 1980. Observations on length-weight relationships and distribution of the red spot king prawn, *Penaeus longistylus* Kubo, in Western Australian waters. *Aust. J. Mar. Freshwater Res.* **31**: 547-552.
- PENN, J.W. and N.G. HALL. 1974. Morphometric data relating to commercial prawns from Shark Bay, Western Australia. *Fish. Res. Bull. West. Aust.* No. 15.
- RAMAMURTHY, S. and M. MANICKARAJA. 1978. Relation between tail and total lengths and total and carapace lengths for three commercial species of Penaeid prawns in India. *Indian J. Fish.* **25**(2): 233-237.
- RHODES, G.P. and D.M. HOLDICH. 1984. Length-weight relationship, muscle production and proximate composition of freshwater crayfish *Austropotamobius pallipes* (Lereboullet). *Aquaculture* **37**: 107-123.
- SORAL, R.R. and F.J. ROHLF. 1981. *Biometry*. 2nd edn. San Francisco: Freeman.

(Received 10 June 1993)